



# Anterior Cervical Disc Replacement/ Arthroplasty (ACDR/ACDA) Post Op Instructions

Kyle Kesler, MD

(423) 624-2696 | [Sportmed.com](http://Sportmed.com)

---

**Anterior Cervical Disc Replacement/Arthroplasty (ACDR/ACDA)** removes disc or bone spurs that are putting pressure on the spinal cord and/or nerve roots. A prosthetic, or implant, is inserted where the disc was removed.

**Post-op pain:** Usually mild to moderate (not severe).

**Goals of Surgery:**

1. Decrease neck and arm pain
2. Stop symptoms of spinal cord compression from getting worse

**Top 4 things your surgeon wants you to know:**

1. Pain in the back of the neck and between the shoulder blades is common after an ACDR/ACDA. It is also normal to have some swallowing difficulty. This usually gets better over the next few weeks. If you have trouble breathing after surgery, call 911 or go to the Emergency Room immediately.
2. You should avoid Nicotine before and after your surgery to help your healing.
3. You need to take Celebrex for two weeks. You will need to take this even if no pain is present. This helps prevent unwanted bone growth called heterotopic ossification.
4. In most cases, after an ACDA, you do not need a collar. If your surgeon did give you one based on your condition, you should wear your collar as directed by your surgeon until your first post-operative appointment. **It is okay to move your neck after surgery.**

## Answers to Frequently Asked Questions about your ACDR/ACDA

### Airway/Breathing/Throat Pain

- If food sticks when you eat, it is called “dysphagia.” This is due to swelling around your surgery site and will most likely resolve in a few weeks. It is normal to feel “tight” for up to 6 weeks, but you should be able to advance your diet slowly and this does improve with time.
- Stick with soft foods that are easy to swallow. Take small bites and chew your food well. Advance your diet to normal as tolerated.
- Having a sore throat is normal, this gets better over the next few weeks after surgery.
- Ice and popsicles help with sore throat.
- If you have **any trouble breathing or have excessive swelling in your neck**, call 911 or go to the emergency room immediately.

### Dressing/Shower

- You can take your dressing off 1-2 days after surgery. **You can take a shower 2-3 days after surgery, once the incision is sealed and not open or leaking fluid.**
- Ok for gentle soap and water to run over the incision, do not scrub, pat dry with a towel. Please avoid tub baths, swimming pools and hot tubs until the incision is completely healed (4-6 weeks).
- **Occasionally your incision may bleed and your dressing may get saturated with blood, this is okay.** The dressing can be changed to a new, clean dressing.

### Incision care

- If there is no drainage, your incision can be left open to air without a dressing after 3 days.
- If there is drainage, cover with a clean and dry dressing. If it does not slow down or stop after a few days, you may need to call the office to have your wound checked.

### Pain/Weakness

- **Surgical neck pain and muscle spasms are normal after a spine surgery.** This usually gets better over the next few weeks.
- Numbness, tingling and weakness that you had before surgery may take time to improve.
- If you develop significant new weakness after you get home, you should call the office: (423)-624-2696 or go to the Emergency Room.

### Nicotine/smoking

- You should avoid Nicotine before and after your surgery to help your healing.

## Collars

- **Not everyone gets/needs a collar after surgery.**
- In most cases, after an ACDA, you do not need a collar. If your surgeon did give you one based on your condition, you should wear your collar as directed by your surgeon until your first post-operative appointment. **It is okay to move your neck after surgery.**
- **It is ok to take the collar off while showering/eating/hygiene purposes.**

## Anti-inflammatories (NSAIDS)

- You need to take Celebrex for two weeks. This helps prevent unwanted bone growth called heterotopic ossification. You are able to take NSAIDS postoperatively

## Pain Medication/Refills

- If you need refills on your prescriptions, please contact CSMO 2-3 days before you are out of medications so we have sufficient time to process your request. **Refill requests on Friday afternoons/holidays will likely be addressed on the next business day.**
- **Opioids are an addictive medication, therefore, you should start weaning off opioid pain medications on your own as soon as you are able to (Ex: Hydrocodone/Oxycodone).**

## Other Medications

- Tylenol (Acetaminophen): Your pain medication likely has acetaminophen in it. Taking additional Tylenol/acetaminophen can put you over the daily recommended 3,000mg, which can harm your liver. Do not exceed this amount.
- Muscle Relaxers: One of the side effects of your muscle relaxer is drowsiness. If you begin to feel too drowsy and you are not able to get up safely to ambulate, decrease the frequency of your muscle relaxer.

## Activity

- When you go home you may get up and walk... **we want you to be active!!**
- You may go up and down stairs, but make sure to hold on to the rail and have someone with you.
- **You should avoid excessive bending and twisting of your neck/back and may not lift anything over 10 pounds until cleared by your surgeon - typically 6 weeks post- op.**

## Driving

- No driving until you are off of all narcotic/sedative medications AND can move well enough to be safe behind the wheel, **this is usually after your first post-operative appointment.**
- **Check with your doctor at your first follow up appointment (4-6 weeks post op) about when you should start driving.**

### **Constipation/Bloating:**

- **A common side effect of narcotic pain medication is constipation.**
- Taking over the counter stool softeners/laxatives may help. Please follow package instructions.
- Stool softeners/Laxatives include:
  - Milk of Magnesia, Miralax, Dulcolax suppository, fleets enema, Magnesium Citrate.Drinking fluid, activity, and diets high in fiber are also helpful in relieving constipation.

### **Follow up with Primary Care Provider**

- If you have any of the below problems, **we suggest you see your Primary Care Provider within 1 week after your surgery** to make sure your other medical issues are doing ok.
- **See your primary care provider if you have a history of:** Heart problems, lung problems, stroke, diabetes, are over the age of 65, are taking a blood thinner, or have several medical problems, or take greater than 10 prescription medications.

## **Emergency Room Use**

**You have trouble breathing, chest pain, or significant NEW weakness after your surgery, please go to a CHI Memorial Emergency Room immediately.**

**Contact our office during office hours : 8:00 am - 4:30 pm at (423) 624-2696.**

**(800) 757-2696 (outside Chattanooga)**

**After office hours phone calls go to on call surgeon/PA. Please only call after hours in urgent situations.**

**If you have any other problems related to your surgery, PLEASE CALL OUR OFFICE BEFORE GOING TO THE EMERGENCY ROOM, as we can likely address your problem or get you an appointment scheduled and save you a trip to the ER.**

**Thank you for choosing Center for Sports Medicine and Orthopedics.**